

Math 350
Spring, 2000

HOMWORK #4

Do 50 points of the following problems (due 2/10/00).

- 15 pts. 1 Find the $n, M,$ and d of the binary code whose generator matrix is

$$\begin{pmatrix} 1 & 0 & 0 & 1 & 1 & 0 & 1 \\ 0 & 1 & 0 & 1 & 0 & 1 & 1 \\ 0 & 0 & 1 & 0 & 1 & 1 & 1 \end{pmatrix}$$

Since the rows are linearly independent, this code has $M = 8$; the length is $n = 7$. The codewords in this code will have even weight since the sum of any two rows has even weight, and it is not possible to have a word of weight 2, so the minimum weight is $d = 4$.

More answers on next pages!

15 pts. **2** Put the following generator matrix into standard form.

$$\begin{pmatrix} 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 \\ 0 & 1 & 0 & 1 & 0 & 1 & 0 & 1 \\ 0 & 0 & 1 & 1 & 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 0 & 1 & 1 & 1 & 1 \end{pmatrix}$$

$$\begin{pmatrix} 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 \\ 0 & 1 & 0 & 1 & 0 & 1 & 0 & 1 \\ 0 & 0 & 1 & 1 & 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 0 & 1 & 1 & 1 & 1 \end{pmatrix}$$

\mathbb{R}

$$\begin{pmatrix} 1 & 0 & 1 & 0 & 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 1 & 0 & 1 & 0 & 1 \\ 0 & 0 & 1 & 1 & 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 0 & 1 & 1 & 1 & 1 \end{pmatrix}$$

\mathbb{R}

$$\begin{pmatrix} 1 & 0 & 0 & 1 & 1 & 0 & 0 & 1 \\ 0 & 1 & 0 & 1 & 0 & 1 & 0 & 1 \\ 0 & 0 & 1 & 1 & 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 0 & 1 & 1 & 1 & 1 \end{pmatrix}$$

\mathbb{R}

$$\begin{pmatrix} 1 & 0 & 0 & 1 & 0 & 1 & 1 & 0 \\ 0 & 1 & 0 & 1 & 0 & 1 & 0 & 1 \\ 0 & 0 & 1 & 1 & 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 0 & 1 & 1 & 1 & 1 \end{pmatrix}$$

\mathbb{R}

$$\begin{pmatrix} 1 & 0 & 0 & 0 & 1 & 1 & 1 & 0 \\ 0 & 1 & 0 & 0 & 1 & 1 & 0 & 1 \\ 0 & 0 & 1 & 0 & 1 & 0 & 1 & 1 \\ 0 & 0 & 0 & 1 & 0 & 1 & 1 & 1 \end{pmatrix}$$

- 20 pts. **3** Let C be a code. Define $C^\perp = \{x \in F_2^n \mid \sum_{i=1}^n x_i c_i = 0 \pmod{2} \text{ for every } c \in C\}$. Find E_n^\perp , and argue why you think that you have the correct answer.

$E_n^\perp = \{000 \cdots 0, 111 \cdots 1\}$. Certainly, these two codewords are contained in E_n^\perp . There are two ways to show that this is it: (1) If E_n^\perp contained any word e so that $1 \leq w(e) \leq n - 1$, then e has at least one position with a 1 and one position with a 0. We form a codeword in E_n so that the codeword has a 0 in every position except one of the positions with a 1 and one with a 0 (the weight of this new codeword is 2). When we take the dot product of e with this new codeword, we get 1, contradicting the definition. Thus, e could not have been in E_n^\perp . (2) The dimension of E_n is $n - 1$, so the dimension of its \perp is $n - (n - 1) = 1$. Since we know that $\{000 \cdots 0, 111 \cdots 1\}$ is contained in E_n^\perp , it must be all of it.

- ★ 30 pts. **4** From the vector space $V(2, q)$, an incidence structure A_q is defined as follows. The 'points' of A_q are the vectors of $V(2, q)$. The 'lines' of A_q are the one-dimensional subspaces of $V(2, q)$ and their cosets. The point P 'belongs to' the line L if and only if P is in L . Prove that A_q is a $(q^2, q, 1, q + 1, q^2 + q)$ design. List the points and lines of A_2 and A_3 .

Clearly there are q^2 points (there are that many vectors in $V(2, q)$). Each one-dimensional subspace is all the scalar multiples of a single vector, and there are q scalars, so there are that many points on a block. Take any pair of points: if they are scalar multiples, then they will be on the line determined by either of the (nonzero) points. If they are not scalar multiples, then if we subtract them, they are both on the line $p_1 + \langle p_2 - p_1 \rangle$. In either case, this is the only line they will be a part of. To see that each point is on $q + 1$ lines, take any point p and any line l not containing p . There will be a distinct line through p and each of the q points on l , and there will be another line through p that is a coset of l that is parallel to l . That makes $q + 1$. To count the number of lines, we need to count the number of one dimensional subspaces (there are $\frac{q^2-1}{q-1} = q + 1$ of them) and multiply that number by the number of cosets of each line (there are q) for a total of $q^2 + q$ lines.

The points of A_2 are 00,01, 10, and 11. The lines are $\{00, 01\}$; $\{10, 11\}$; $\{00, 10\}$; $\{01, 11\}$; $\{00, 11\}$; and $\{01, 10\}$. The points of A_3 are 00,01,02, 10,11,12,20,21, and 22. The lines are $\{00, 01, 02\}$; $\{10, 11, 12\}$; $\{20, 21, 22\}$; $\{00, 10, 20\}$; $\{01, 11, 21\}$; $\{02, 12, 22\}$; $\{00, 11, 22\}$; $\{10, 21, 02\}$; $\{20, 01, 12\}$; $\{00, 12, 21\}$; $\{10, 22, 01\}$; and $\{20, 02, 11\}$.